

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>4</sup> : <b>A61K 37/02</b>	<b>A2</b>	(11) International Publication Number: <b>WO 88/ 00833</b> (43) International Publication Date: 11 February 1988 (11.02.88)
--	-----------	--

(21) International Application Number: PCT/US87/01994

(22) International Filing Date: 7 August 1987 (07.08.87)

(31) Priority Application Numbers: 894,910  
076,625(32) Priority Dates: 8 August 1986 (08.08.86)  
23 July 1987 (23.07.87)

(33) Priority Country: US

(71) Applicant: BOARD OF REGENTS, THE UNIVERSITY OF TEXAS SYSTEM [US/US]; 201 West Seventh Street, Austin, TX 78701 (US).

(72) Inventor: LIPTON, James, M. ; 100662 Royal Springs Drive, Dallas, TX 75229 (US).

(74) Agent: PARKER, David, L.; Arnold, White &amp; Durkee, Houston, TX 77210 (US).

(81) Designated States: AT, AT (European patent), AU, BB, BE (European patent), BG, BJ (OAPI patent), BR, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CM (OAPI patent), DE, DE (European patent), DK, FI, FR (European patent), GA (OAPI patent), GB, GB (European patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL, NL (European patent), NO, RO, SD, SE, SE (European patent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent).

**Published***Without international search report and to be republished upon receipt of that report.*

(54) Title: ANTIPYRETIC AND ANTI-INFLAMMATORY PEPTIDES

## (57) Abstract

An antipyretic tripeptide, having the amino acid sequence lysine-proline-valine, and a method for utilizing the tripeptide to reduce fever and inflammation in mammals. The tripeptide can either be isolated from natural sources or chemically synthesized. A "protected" tripeptide having greater antipyretic potency and duration of action is also disclosed. The "protected" tripeptide contains an acyl group, such as an acetyl or a dibenzyl oxy carboxyl group, at its amino terminals and is amidated or esterified at its carboxyl terminals. Further, improved antipyretic potency and direction of action can be achieved through the co-administration of copper salts with the tripeptide.

***FOR THE PURPOSES OF INFORMATION ONLY***

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

-|-

5

10

ANTIPYRETIC AND ANTI-INFLAMMATORY PEPTIDES

15

This invention relates to a new pharmaceutical composition useful for the treatment of pyrexia and inflammation. More particularly, this invention relates to a tripeptide sequence contained in alpha-Melanocyte Stimulating Hormone and ACTH which has been identified as an antipyretic and anti-inflammatory agent.

25 There are two classes of agents presently in common usage as antipyretic agents, the salicylates and the para-aminophenol derivatives. The salicylates, characterized by acetylsalicylic acid (i.e., aspirin), are the most extensively employed antipyretic agents. Aspirin is the prototype for both the salicylates and other drugs with similar effects and is the standard of reference for comparison and evaluation of these agents. The antipyretic effect of aspirin is usually rapid and effective in febrile patients. The salicylates act to reset the

35 "thermostat" for normal temperature.

Although aspirin is generally well-tolerated by most individuals, a number of toxic side effects are associated with its use. Of particular concern is salicylate-induced gastric ulceration and sometimes hemorrhage. Exacerbation  
5 of peptic ulcer symptoms (heartburn, dyspepsia), gastro-intestinal hemorrhage, and erosive gastritis have all been reported in patients taking aspirin. Other less common side effects include tinnitus and hearing loss, changes in acid-base balance and electrolyte pattern, and respiratory  
10 alkalosis. Although generally such side effects are not particularly dangerous, they tend to reduce patient compliance. Other salicylate derivatives, which are generally employed for their analgesic and/or anti-inflammatory activity, demonstrate increased toxicities  
15 relative to aspirin.

The para-aminophenol derivatives, acetaminophen and phenacetin, are alternatives to aspirin for its analgesic and antipyretic uses. Acetaminophen has somewhat less  
20 overall toxicity and is generally preferred over phenacetin. Because acetaminophen is well-tolerated and lacks many of the undesired effects of aspirin, it has been gaining favor as the "common household analgesic." However, its suitability for this purpose is questionable: in  
25 acute overdosage, acetaminophen can cause fatal hepatic necrosis. In addition, phenacetin may cause methemoglobinemia and hemolytic anemia as a form of acute toxicity, but more commonly as a consequence of chronic overdosage. These agents are about equipotent with as-  
30 pirin in the treatment of pyrexia.

Recent advances in the study of alpha-Melanocyte Stimulating Hormone (hereinafter referred to as "MSH") have demonstrated that this protein is active in the  
35 treatment of pyrexia. Alpha MSH is a 13-amino acid peptide derived from Adrenocorticotrophic Hormone ("ACTH").

Both MSH and ACTH share a 13-amino acid sequence that is effective in modulating body temperature. There is evidence that these neuropeptides can influence centrally mediated processes, including central control of body temperature. Both peptides lower core temperature of afebrile rabbits when given peripherally or centrally in sufficient dosages. Much smaller dosages reduce fever without altering normal temperature.

- 10        Alpha MSH is found in brain regions that govern temperature regulation, including the anterior hypothalamus and the septum. The concentration of alpha MSH in the septum rises during fever, and the concentration in the arcuate nucleus tends to decline at the same time.
- 15        Studies comparing the antipyretic activity of centrally-administered alpha MSH to the widely-used antipyretic, acetaminophen indicate that alpha MSH is much more potent in reducing fever than acetaminophen, and that alpha MSH was more than 2500 times more potent by weight than
- 20        acetaminophen in reducing fever. No endogenous substance other than ACTH is known to have such potency in reducing fever.

      The antipyretic potency of alpha MSH and the fact that this peptide reduces fever even when given peripherally may have clinical significance. ACTH was used to reduce clinical and experimental fever soon after it was first described, but this peptide also stimulates corticosteroid release, and can, with repeated administration, result in Cushing's syndrome. On the other hand, the shorter alpha MSH molecule, which is derived from ACTH, does not stimulate steroid release and there appears to be no irreversible deleterious effects when given to rabbits or to man.

With respect to ACTH (amino acids 1-39 of proopio-cortin), it has previously been known that due to its corticosteroid stimulatory effect, this protein was active in the treatment of inflammation. However, shorter ACTH-related peptides such as alpha-MSH (amino acids 1-13 of ACTH) which do not exhibit corticotropic activity have not been shown to have anti-inflammatory action, and there has previously been no basis to suggest such a role for peptides which correspond to amino-terminal portions of ACTH yet which exhibit no corticotropic activity.

The present invention provides a pharmaceutical composition useful in the treatment of pyrexia and inflammation. The active component of this pharmaceutical composition is a peptide which includes an amino acid sequence corresponding to amino acids 11 through 13 of alpha MSH, lysine-proline-valine ("lys-pro-val"). In its most general scope, the invention is directed to peptides of from 3 to 13 amino acids, which peptides have sequences corresponding to that of alpha-MSH and include at least the lys-pro-val sequence thereof. In more particular embodiments, the invention is directed to the tripeptide itself.

Preferably, the tripeptide itself is administered to achieve maximal benefits in accordance herewith, preferably in a biologically "protected" form. When the tripeptide is "protected" through acylation of the amino terminus and/or amidation of the carboxyl terminus, the resulting tripeptide demonstrates an increase in pharmacologic activity. Similarly, when the tripeptide, whether "protected" or "unprotected", is co-administered with copper ion, a further increase in antipyretic activity is observed.

The present invention provides a method for treating pyrexia and/or inflammation in an individual in need of such treatment in which an effective dose of a peptide which includes the tripeptide sequence is administered to the pyretic individual. Moreover, such peptides may be used in the treatment of both generalized or localized inflammation and, therefore, is a useful alternative to steroidal and salicylate anti-inflammatory agents. Anti-inflammatory activity is observed following administration of the tripeptide to animals at doses approximately equal or greater than those used to demonstrate antipyresis, and testing the reactivity of treated and control animals to inflammatory challenge. Therefore, the peptides of the present invention may be used both as an antipyretic and as an anti-inflammatory agent when administered at a selected dose to a patient in need.

In particular embodiments herein, doses of the tripeptide effective for the expression of anti-inflammatory activity, for example, for the reduction of inflammation-associated swelling and/or capillary permeability, are shown to be roughly similar on a weight basis to those of hydrocortisone, an accepted standard for anti-inflammatory activity. The sensitive "skin-blueing" assay was employed to test the ability of the tripeptide to inhibit the capillary permeabilizing effects of inflammatory agents such as histamine. In this assay, the protected Lys-Pro-Val tripeptide exhibited antihistamine effects, and in particular, a reduction in histamine-mediated increases in capillary permeability, at intravenous dosages as low as 1.25 ug protected tripeptide/kg body weight. Moreover, in the traditional carrageenan/rat paw edema test, intraperitoneally-administered tripeptide demonstrated an ability to inhibit carrageenan-induced swelling of rat paws on a per weight basis commensurate with hydrocortisone. Accordingly, from such observations



it can be concluded that the tripeptide lys-pro-val is an effective anti-inflammatory when administered to a patient at a dose ranging from as low as 1 to 10 ug/kilogram, to preferred ranges of on the order of about 0.2 to about 3 mg/kg/day.

Figure 1. Amino acid sequence of alpha-MSH

Figure 2. Comparison of inhibition of paw swelling caused by Ac-Lys-Pro-Val-NH<sub>2</sub> (100 mg/kg) and hydrocortisone (100 mg/kg) in rats. Each score is the mean % change in paw volume relative to the change in matched control paw volume.

Peptides used in the practice of the present invention include the sequence lysine-proline-valine. This tripeptide sequence is characterized as follows.

In its naturally occurring form, the tripeptide sequence (lysine-proline-valine) comprises amino acid numbers 11-13 of alpha-Melanocyte Stimulating Hormone (hereinafter "MSH") and ACTH. This finding may explain the antipyretic activity of the amino-terminal portion of ACTH (amino acid numbers 1-24 of proopiomelanocortin) and MSH (amino acid numbers 1-13 of ACTH and proopiomelanocortin; see Figure 1), both of which exhibit the tripeptide sequence within their structure. Therefore, both MSH and ACTH represent potential naturally occurring sources from which the antipyretic tripeptide can be obtained, or, in the case of alpha-MSH, which can be used directly in accordance with less preferred embodiments.

Due to the high corticotropic effect of ACTH, which can lead to incidences of Cushing's Syndrome, the invention is generally directed to the alpha-MSH sequence and peptides thereof, so long as such peptides include at

least the lys-pro-val sequence. This is based on the finding that alpha-MSH (amino acids 1-13 of ACTH) does not exhibit the corticotropic effect of ACTH and, instead, appear to exert their anti-inflammatory action directly rather than through a corticosteroid intermediate.

In preferred embodiments, the tripeptide can be isolated from MSH. This can be accomplished by first fragmenting the MSH protein into four smaller peptides through total digestion with the proteolytic enzyme, chymotrypsin. Paper electrophoresis of the digestion products reveals four major products, one of which is the tetrapeptide, glycine-lysine-proline-valine, which may be used directly. However, the glycine residue can be removed by partial acid hydrolysis to yield the tripeptide.

Peptides in accordance with the invention can also be obtained by chemical synthesis. This is accomplished by way of peptide bond formation between the appropriate amino acids. Amino acids are amphoteric molecules which contain both an acidic ( $-\text{COOH}$ ) moiety and a weakly basic ( $-\text{NH}_2$ ) moiety. Peptide bond formation ( $-\text{CONH}-$ ), therefore, is accomplished through a nucleophilic attack of the amine group on the carboxylic function.

In forming a peptide bond between two hypothetical amino acids, X and Y, four possible dipeptides may be produced: X-Y, Y-X, Y-Y, and X-X. Therefore, in order to reduce the possible structures that may be formed in such an interaction, the amino or carboxy terminus of the appropriate amino acid must be first "protected" so as to preclude a reaction involving the "protected" moiety. For example, if "c" represents a protected carboxy terminus and "n" a protected amino terminus, then an interaction

involving cX and Yn could generate only one structure, cX-Yn.

5        However, in order to be chemically useful for synthetic purposes, the protecting groups must be removable. In general, carboxy groups can be protected by esterification or amidation of the -COOH to -COO-alkyl or -CONH<sub>2</sub>. The preferred alkyl groups for the carboxy terminus include methyl and benzyl residues, yet other alkyl groups, such  
10 as ethyl, propyl, butyl, p-nitrobenzyl or p-methoxybenzyl groups, can be utilized.

Likewise, the amino terminus is protected by acylation, introducing a carboxyl group such as an acetyl



compared. From such assays, it was found that approximately equal weights of the protected tripeptide (diacetyl-Lys-Pro-Val-NH<sub>2</sub>) and hydrocortisone resulted in a roughly equivalent overall response in reducing the degree of carrageenan-induced swelling.

Based on the foregoing and additional observations, it is found that dosages on the order of 0.2 to about 3 mg of protected tripeptide per kilogram body weight per day will result in advantages in accordance with the present invention in terms of effective anti-inflammatory action. Generally, it will be preferred to administer doses of on the order of 0.35 and about 1.5 mg of protected tripeptide/kg body weight/day to achieve the greatest degree of anti-inflammatory benefit. These dose ranges are derived from the aforementioned observation of approximately equipotency of the tripeptide and hydrocortisone, and the general knowledge in the art regarding effective dose ranges of hydrocortisone (see, e.g., Goodman et al. (1985), The Pharmacological Basis of Therapeutics, 7th Edition).

In accordance with the invention, it will generally be preferred to administer the tripeptide parenterally, for example, intramuscularly or intravenously. However, due to its small size, membrane permeability and relatively acid-stable structure, it will be recognized that the tripeptide may be administered orally, albeit at somewhat higher doses. In this regard, it is believed that doses of on the order of about 0.2 to about 3.5 mg/kg/day will achieve benefits in accordance herewith.

Pharmaceutical preparations of the tripeptide, preferably a protected tripeptide such as diacetyl-Lys-Pro-Val-NH<sub>2</sub>, comprise generally the active agent in combination with pharmaceutically acceptable buffers,

diluents, stabilizers and the like. For a fairly complete listing of various techniques, including a variety of agents and additives useful in the preparation of acceptable pharmaceutical compositions, one may wish to refer to 5 Remington's Pharmaceutical Sciences, 16th ed., 1980, Mack Publishing Co., incorporated herein by reference.

In a preferred pharmaceutical composition, approximately 100 to 500 mg of diacety-Lys-Pro-Val-NH<sub>2</sub> is 10 dispersed in about 1 to 7 cc. of sterile isotonic saline, including a pharmacologically accepted buffer to maintain a pH of about neutral. For intravenous administration, for example, to a patient suffering from arthritis, or severe allergic reaction or various other diseases 15 involving inflammatory processes, it will generally be desirable to administer about 0.2 to about 3.5 mg/kg/day of the agent by slow infusion over a period of time (up to several hours). Where infusion is impractical, the agent is administered in the form of an intramuscular injection, 20 preferably in combination with a lipophilic carrier and at

the protein. As displayed in the table below, amino acids are assigned a hydropathic index on the basis of their hydrophobicity and charge characteristics. It is believed that the relative hydropathic character of the amino acid  
5 determines the secondary structure of the resultant protein, which in turn defines the interaction of the protein with its receptor.

In the case of the present peptides, it is believed  
10 that biological functional equivalents may be obtained by substitution of amino acids having similar hydropathic values. As used herein, a biological functional equivalent is defined as a protein that is functionally equivalent in terms of biological functional equivalent is  
15 defined as a protein that is functionally equivalent in terms of biological activity. Thus, for example, isoleucine of leucine have a hydropathic index of +4.5 and +3.8, respectively, can be substituted for valine (+4.2), and still obtain a protein having like biological  
20 activity. Alternatively, at the other end of the scale, lysine (-3.9) can be substituted with arginine (-4.5), and so on. In general, it is believed that amino acids can be successfully substituted where such amino acid has a hydropathic score of within about +/- 1 hydropathic index  
25 unit of the replaced amino acid.

Amino acid

Hydropathic Index



desired (a biologically functional equivalent of lys-pro-val), it will be apparent that dibenzylloxycarbonyl-conjugated arginine ("Z-arg") should be employed in the place of "Z-arg"). Moreover, where, for example, gly-lys-pro-val is desired, it will be apparent that "Z-gly" should be employed as the starting reagent and synthetic steps employed as set forth to sequentially add the lys, pro and val residues, respectively. These and all other modifications to achieve the various peptides are well known and will be apparent to those of skill.

EXAMPLE I: CHEMICAL SYNTHESIS OF L-LYSINE-L-PROLINE-L-VALINE

The tripeptide was custom synthesized by Bachem, Inc., Torrance, Ca., as follows:

1. Z-Lys-Pro-OMe Preparation

50 mmoles (20.7 grams) of dibenzylloxycarbonyl-conjugated lysine ("Z-lys") in 200 ml of methylene chloride was combined with 50 mmoles (8.3 grams) of proline methyl ester (pro-OMe) in 100 ml dimethyl formamide. The mixture was added to a conical flask and cooled to -5°C with stirring. 50 mmoles (5.5 ml) of N-methyl morpholine was added, followed by the addition of 10.3 grams of dicyclohexyl-carbodiimide in 20 ml of methylene chloride and the reaction mixture was stirred overnight. The mixture was then filtered from urea and the filtrate concentrated in vacuo. The residue was taken up in ethyl acetate and washed successively with sodium bicarbonate solution, water, 1N hydrochloric acid, and water. Ethyl acetate was removed in vacuo and the oily product was saponified without purification.

2. Removal of the -OMe Carboxy Terminus  
"Protecting" Group.

The oily product from the previous experiment was  
5 dissolved in methanol (200 ml) and treated with 2N sodium  
hydroxide (25ml) for an hour. Methanol was removed under  
reduced pressure and the residue was taken up in water and  
acidified with 6N hydrochloric acid. The product was  
10 extracted with ethyl acetate, and the organic layer was  
washed with water and dried over sodium sulphate. Ethyl  
acetate was removed in vacuo and the residue was tri-  
tured with hexane. The product was checked by thin  
layer chromatography using chloroform: methanol: acetic  
acid (95:4:1).

15

3. Preparation of Z-Lys-Pro-Val-OBe

The next step in the synthesis of the tripeptide  
involved the addition of a carboxy-protected valine resi-  
20 due (Val-OBe). The protecting group in this instance was  
a benzyl ester.

37 mmoles (19 grams) of Z-Lys-Pro obtained from step  
2 above was dissolved in 200 ml distilled tetrahydrofuran.  
25 This solution and 4.1 ml N-methyl morpholine were mixed  
together and cooled to -15°C with stirring. Isobutyl-  
chloroformate (5)(ml) was added and the mixture stirred  
for 5 minutes at -10°C. Concurrent with the above, 35  
mmoles (13.2 grams) of valine benzyl ester tosylate was  
30 dissolved in 100 ml dimethyl formamide. The mixture was  
cooled to -10°C and neutralized with N-methyl morpholine  
(4ml). This was added to the above mixed anhydride and  
stirred overnight. The mixture was then filtered from  
urea and the filtrate concentrated in vacuo. The residue  
35 was taken up in ethyl acetate and washed successively with  
sodium bicarbonate solution, water, 1N hydrochloric acid,

and water. The crude product was purified on a silica gel column using chloroform-methanol (95:5). Fractions containing the pure product were determined by thin layer chromatography utilizing the same solvent as for step 2.

5 The appropriate fractions were pooled.

#### 4. Removal of the Protecting Groups Z- and -OBe

9 grams of the protected tripeptide produced in step  
10 3 above was hydrogenated in an acetic acid-water-methanol mixture in the presence of Pd/BaSO<sub>4</sub> overnight. It was filtered from the catalyst and the filtrate was evaporated in vacuo to give an oily residue. This was triturated with absolute ethanol and absolute ether to yield 3 grams  
15 of the crystalline product. The product was checked by thin-layer chromatography using a solvent system composed of butanol: acetic acid: water: pyridine (20:6:11:24).

#### 20 Chemical Synthesis of diacetyl-L-Lysyl-L-Prolyl-L-Valyl-NH<sub>2</sub>

The protected tripeptide, diacetyl-L-Lysyl-L-Prolyl-L-Valyl-NH<sub>2</sub> can also be prepared by the chemical techniques described above in steps 1-3. For example, in step  
25 1, the starting material would be diacetyl conjugated lysine. In step 3, the valine-benzylester is substituted with valyl-amide.

#### 30 EXAMPLE II: ANTIPYRETIC ACTIVITY OF L-LYS-L-PRO-L-VAL

##### Production of Leukocytic Pyrogen

Leukocytic pyrogen is a molecule capable of producing  
35 transient fever in mammals which is produced by incubating rabbit leukocytes with Salmonella typhosa endotoxin. More

specifically, to produce leukocytic pyrogen, donor rabbits were first sacrificed by decapitation. Blood was collected in a heparinized pyrogen-free beaker. Heparinized 50-ml glass centrifuge tubes were filled 3/4 full with whole blood, saline was added to fill each tube, and the solution was gently mixed. The tubes were then centrifuged at low speed for 20 min. The buffy coat was drawn off and placed in pyrogen-free flasks. Lactate Ringer's solution equal in volume to one-half that of the red cell layer, was added along with Salmonella typhosa endotoxin (Difco, No. 0901) also in Ringer's solution (1 ug/ml), to the buffy coat. The mixture was incubated at 38°C in a shaking water bath for 4 hours. The solution was centrifuged, filtered (Nalgene, 0.20 microns), and the leukocytic pyrogen-containing filtrate was stored at 4°C. Samples of leukocytic pyrogen were heated to 70°C for 2 hours and injected intravenously to test for endotoxin contamination. Only characteristic leukocytic pyrogen fevers occurred and no prolonged fevers were observed, indicating that the leukocytic pyrogen was free of endotoxin and other heat-stable pyrogens.

Injectons were 50 ul in volume and were followed by a 20 ul saline flush. Intravenous injections were made via the marginal ear vein. Intravenous leukocytic pyrogen injections were 0.07 ml of a stock solution made up of a mixture of leukocytic pyrogen derived from 4 donors. When injected, leukocytic pyrogen stock solution was diluted with nonpyrogenic isotonic saline. Injections were made with commercial nonpyrogenic syringes. Glassware was washed with chromic acid, rinsed with deionized water and heated to 200°C for a minimum of 2 hours to insure that it was pyrogen free.

### Animal Procedures

Adult New Zealand white rabbits were used for the development of an antipyretic assay. The rabbits were  
5 housed individually in a 21-23°C environment with a 12 hour light/dark cycle; food and water were available ad libitum. Central nervous system injections of the antipyretic agents were performed as follows: The animals were pretreated with ketamine hydrochloride and promazine  
10 (Ketaset Plus, Bristol Labs, 0.4 ml/kg. intramuscularly) and anesthesia was induced and maintained by inhalation of methoxyfluorane (Metafane, Pitman-Moore, Inc.) and an N<sub>2</sub>O-O<sub>2</sub> mixture. Rabbits were placed in a Kopf rabbit stereotaxic instrument and a stainless steel cannula (No.  
15 201, David Kopf Instruments) was inserted into a lateral ventricle at a point 1.0 mm anterior to the bregma and 2.7 mm lateral to the midline. The cannula was lowered until cerebrospinal fluid appeared inside the well of the cannula. Stainless steel screws and dental acrylic were  
20 used to anchor the cannula to the calvarium. Benzathine penicillin G (Bicillin, Wyeth Laboratories) was given post-operatively (150,000 units intramuscularly).

Experimental rabbits that were to be used in the  
25 antipyresis study were restrained in conventional holders and a thermistor probe (Yellow Springs International, No. 701) was inserted about 100 mm into the rectum and taped to the tail. In certain experiments, another thermistor probe (Yellow Springs International, No. 709) was attached  
30 to the dorsal surface of the ear. Temperature recordings were made every 10 minutes via a MINC 11 online computer connected to a digital temperature recorder (Datalogger, United Systems Corp.) At least 1 hour was allowed after the probes were inserted before injections were made, and  
35 all experiments were separated by at least 48 hours. Experiments were run in an environmental chamber at 23°C.

The average thermal response, the mean change in temperature ( $^{\circ}\text{C}$ ) over the duration of the response measured in hours, was calculated for each response and the paired t-test was used for statistical analysis of the data. The time period over which the experimental antipyretic temperature response was determined was generally set by the duration of the control response for the individual animal. The control response begins with the first deviation from baseline temperature and continues until the temperature returns to baseline or to the point nearest the baseline within 5 hours. The mean change in temperature for each 10 minute intervals during this time period are summed and divided by the total number of 10 minute periods.

15

#### Assay Protocol and Results

Immediately after the tripeptide, L-Lys-L-Pro-L-Val, was synthesized as described above, it was dissolved in sterile non-pyrogenic isotonic saline and stored frozen in aliquots until just prior to use. Before any injections of the tripeptide were given, leukocytic pyrogen was tested in each animal in order to establish its sensitivity to the pyrogen and to test for endotoxin activity.

To induce fever in the test animals, 0.15 ml of a stock solution of leukocytic pyrogen was injected into a marginal ear vein. Pyrogen from several batches was used, but each animal received pyrogen from the same batch throughout each series of experiments.

Injections of the tripeptide were given 30 minutes after the pyrogen, into the intraventricular cannula. Centrally administered tripeptide resulted in an observed antipyresis. Decreases in fever, calculated as percent

reduction of the area under the control fever curve over the apparent duration of action of the peptide (1.5 hours), were 24%, 31%, and 48% for the 0.5, 1.0, and 2.0 milligram doses, respectively. Similarly, intravenous administration of 2, 20 and 200 milligrams of the tripeptide reduced fever 34, 27 and 67% , respectively, during the 1.5 hour period after injection of the leukocytic pyrogen. Control saline injections, centrally administered, caused no significant reduction in body temperature. Similarly, when injections of 200 milligrams were given to afebrile rabbits, no reduction in body temperature was observed.

15      **EXAMPLE III: ANTIPYRETIC ACTIVITY  
OF THE PROTECTED TRIPEPTIDE  
DIACETYL-L-LYS-L-PRO-L-VAL-NH<sub>2</sub>**

Central administration of the acetylated and amidated tripeptide diacetyl-L-Lysine-L-Proline-L-Valine-NH<sub>2</sub> resulted in an increase in the observed antipyresis as well as an increase in the duration of action. Fever induced by intravenous administration of leukocytic pyrogen was reduced more than 50% by 0.5 mg of the protected peptide. The duration of action was at least four hours compared to 1.5 hours observed for the unprotected tripeptide (see Example I). Smaller doses of the protected tripeptide resulted in correspondingly lower antipyresis.

30      **EXAMPLE IV: ANTIPYRETIC ACTIVITY  
OF DIACETYL-L-LYS-L-PRO-L-VAL-NH<sub>2</sub>  
AND COPPER ION**

When 0.5 mg of the protected peptide was given centrally and copper ions (1-10 mg of the cupric chloride salt) were given either centrally or peripherally, the antipyretic effect was greatly augmented and hypothermia

developed that was similar to that that has been observed with large doses of the parent alpha-MSH peptide. The protected peptide thus appears to be at least four times more potent than the unprotected tripeptide. The addition  
5 of copper ions, in doses that have no effect on normal temperature, markedly enhanced the antipyretic, and hypothermic, effects of the protected peptide.

EXAMPLE V: ANTI-INFLAMMATORY ACTIVITY

10

The anti-inflammatory activity of the tripeptide was demonstrated through the use of an animal model developed by Sparrow and Wilhelm (1957), J. Physiol., 137:51-65, incorporated herein by reference. This model relies on  
15 the principal that localized, subcutaneous injections of histamine will result in a localized increase in capillary permeability. When the test animal has been pretreated with blue dye intravenously, the localized histamine injections will elicit blue-colored "weals" around the  
20 injection site. Thus, by preadministration of an effective anti-inflammatory agent, the blue color of the histamine-induced weals will be much less pronounced, with the amount of color reduction being dependent on the relative amount and/or potency of the anti-inflammatory  
25 agent used.

Non-moulting New Zealand white rabbits were used for the Sparrow/Wilhelm assay. The skin of the rabbits back was closely clipped 1-2 days previous to the experiment,  
30 but not depilated, and the rabbits were kept warm until tested. Various amounts of the protected tripeptide (diacetyl-L-lys-1-Pro-L-Val-NH<sub>2</sub>) were injected intravenously into an ear vein approximately 15 minutes prior to intravenous injection of blue dye. Control rabbits  
35 received sham injections. Fifteen minutes following injection of the agent or sham, the rabbits received



approximately 30 mg/kg of Pontamine blue dye as a 2.5% solution in 0.45% saline, into an exposed vein.

Immediately following dye injections, histamine was injected intradermally in a 0.10 ml volume (1.25 mg histamine /0.1 ml volume) at several sites on each side of the spine. In all, one vertical row of six injections were made on each side of the spine. The relative intensity of the resultant blue weals were scored by an independent observer 30 minutes after histamine injection. The results are displayed in Table I below.

TABLE I: Anti-inflammatory Activity  
of the Tripeptide

15

20	No. Animals	Tripeptide	
	<u>Tested</u>	<u>Dose +</u>	<u>Result</u>
	3 (2E, 1C)*	5	E lighter than C
25	2 (1E, 1C)	10	E lighter than C
	2 (1E, 1C)	5	E lighter than C
	2 (1E, 1C)	1.25	E lighter than C
	2 (1E, 1C)	0.625	No difference observed

30

\* E = experimental; C = control

+ Dosages in ug of protected tripeptide per kg body weight, administered intravenously.

35

As will be appreciated from the results displayed in Table I, intravenous doses down to 1.25 ug tripeptide per kg body weight resulted in an appreciable reduction in histamine-induced blue weal formation and is thus indicative of an effective anti-inflammatory action. At doses of 5 and 10 ug/kg, the observed response was even more pronounced. Also as will be appreciated, the anti-inflammatory effect of the tripeptide is observed at relatively lower doses as compared to its anti-pyretic effect.

EXAMPLE VI: CARRAGEENAN/RAT PAW EDEMA ASSAY

A second in vivo bioassay for anti-inflammatory activity was conducted in which the action of the tripeptide was compared to that of hydrocortisone. In this assay, the two agents were given at similar doses and tested for their independent ability to inhibit carrageenan-induced swelling of rats paws. This assay, the rat paw edema test, was conducted generally as it is typically performed in the art, for example, as described by Winter et al. (1962), Proc. Soc. Exp. Biol. Med., 111:544 or in U.S. Patent 4,150,137.

Briefly, the assay was performed as follows. Each of twenty-four male Sprague-Dawley rats was assigned to one of four groups: tripeptide treatment and controls (matched according to body weight and initial paw volume), hydrocortisone treatment and matched controls. The volume of the right rear paw of the test and control animals was determined using standard procedures and a mercury displacement volumetric technique. An intraperitoneal injection of the tripeptide (Ac-Lys-Pro-Val-NH<sub>2</sub>, 100 mg/kg, N=6), of hydrocortisone (100 mg/kg, N=6), or saline (matched volume, N=12) was given to each rat. One hour later 0.5 ml of 1% lambda carrageenan in saline solution

was injected into the right rear paw of the animals and the paw volume was again recorded (baseline measure). Thereafter paw volume was measured each hour for four hours. For comparison of the effects of the two treatments, paw volume of experimental animals measured at hourly intervals was expressed as a percentage of the volume change of their respective matched controls.

The results of this experiment is shown in Figure 2. As will be appreciated from this data, except for the first hour when hydrocortisone markedly inhibited swelling ( $p < 0.05$ , Mann-Whitney test), there was no significant difference in the inhibition of paw edema caused by the tripeptide and hydrocortisone ( $p > 0.20$ ). These results indicated that the tripeptide inhibits inflammation as well as the classic anti-inflammatory agent, when given in an equal dose by weight, albeit with a slight difference in time course. Based on the present results and the known effects of hydrocortisone and inflammation in man, it is concluded that the tripeptide Lys-Pro-Val can be used to reduce inflammation in man, in dosage not markedly different from that of hydrocortisone.

25

\* \* \*

The foregoing invention has been described by way of illustration and example and in terms of standard laboratory techniques employed by the applicant. It will be apparent to those skilled in the art that certain changes and modifications of these procedures may be employed without departing from the spirit and scope of the invention. For example, although chemically synthesized tripeptide was utilized to demonstrate its antipyretic activity, it is contemplated by the inventor that tri-

peptide isolated from natural sources will function equally well. Moreover, it will be apparent that administration of any copper salt, whether it be the sulfate, chloride, or some other similar copper salt, should be as  
5 active as the chloride salt in augmenting the activity of the tripeptide. Similarly, although activity was demonstrated using either intravenous or centrally administered tripeptide, orally administered tripeptide, at higher  
10 doses, should be active in reducing fever. It will be apparent to those skilled in the art that these and other modifications and changes are within the scope of the appended claims.


---

**CLAIMS:**

# AMINO ACID SEQUENCE OF ALPHA-MSH

NH<sub>2</sub> - SERINE - TYROSINE - SERINE -  
METHIONINE - GLUTAMIC ACID -  
HISTIDINE - PHENYLALANINE -  
ARGININE - TRYPTAMINE -  
GLYCINE - LYSINE - PROLINE -  
VALINE - COOH

**Fig. 1**




**Fig. 2**



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>4</sup> : <b>A61K 37/02</b>		A3	(11) International Publication Number: <b>WO 88/ 00833</b>
			(43) International Publication Date: 11 February 1988 (11.02.88)
(21) International Application Number: PCT/US87/01994 (22) International Filing Date: 7 August 1987 (07.08.87) (31) Priority Application Numbers: 894,910 076,625 (32) Priority Dates: 8 August 1986 (08.08.86) 23 July 1987 (23.07.87) (33) Priority Country: US		(74) Agent: PARKER, David, L.; Arnold, White & Durkee, Houston, TX 77210 (US). (81) Designated States: AT, AT (European patent), AU, BB, BE (European patent), BG, BJ (OAPI patent), BR, CF (OAPI patent), CG (OAPI patent), CH, CH (Eu- ropean patent), CM (OAPI patent), DE, DE (Euro- pean patent), DK, FI, FR (European patent), GA (OAPI patent), GB, GB (European patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (Euro- pean patent), MC, MG, ML (OAPI patent), MR (OA- PI patent), MW, NL, NL (European patent), NO, RO, SD, SE, SE (European patent), SN (OAPI pa- tent), SU, TD (OAPI patent), TG (OAPI patent). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments. (88) Date of publication of the international search report: 19 May 1988 (19.05.88)	
(71) Applicant: BOARD OF REGENTS, THE UNIVERSI- TY OF TEXAS SYSTEM [US/US]; 201 West Seventh Street, Austin, TX 78701 (US). (72) Inventor: LIPTON, James, M. ; 100662 Royal Springs Drive, Dallas, TX 75229 (US).			

(54) Title: ANTIPYRETIC AND ANTI-INFLAMMATORY PEPTIDES



## (57) Abstract

An antipyretic tripeptide, having the amino acid sequence lysine-proline-valine, and a method for utilizing the tripeptide to reduce fever and inflammation in mammals. The tripeptide can either be isolated from natural sources or chemically synthesized. A "protected" tripeptide having greater antipyretic potency and duration of action is also disclosed. The "protected" tripeptide contains an acyl group, such as an acetyl or a dibenzyl oxy carboxyl group, at its amino terminals and is amidated or esterified at its carboxyl terminals. Further, improved antipyretic potency and direction of action can be achieved through the co-administration of copper salts with the tripeptide.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				



# INTERNATIONAL SEARCH REPORT

International Application No PCT/US 87/01994

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) <sup>6</sup> According to International Patent Classification (IPC) or to both National Classification and IPC IPC <sup>4</sup> :            A 61 K 37/02														
<b>II. FIELDS SEARCHED</b> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Minimum Documentation Searched <sup>7</sup></div> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 20%; border-bottom: 1px solid black;">Classification System</td> <td style="border-bottom: 1px solid black;">Classification Symbols</td> </tr> <tr> <td style="padding: 5px;">IPC<sup>4</sup></td> <td style="padding: 5px;">A 61 K 37/00</td> </tr> </table> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>8</sup></div>			Classification System	Classification Symbols	IPC <sup>4</sup>	A 61 K 37/00								
Classification System	Classification Symbols													
IPC <sup>4</sup>	A 61 K 37/00													
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT <sup>9</sup></b> <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 10%; border-bottom: 1px solid black;">Category <sup>10</sup></th> <th style="width: 60%; border-bottom: 1px solid black;">Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup></th> <th style="width: 30%; border-bottom: 1px solid black;">Relevant to Claim No. <sup>13</sup></th> </tr> <tr> <td style="vertical-align: top; padding: 5px;">X,Y</td> <td style="vertical-align: top; padding: 5px;">J. Rheumatol., volume 12, no. 5, October 1985, C.W. Denko et al.: "Effects of peptide hormones in urate crystal inflammation", pages 971-975 see the whole document --</td> <td style="vertical-align: top; text-align: center; padding: 5px;">1-7</td> </tr> <tr> <td style="vertical-align: top; padding: 5px;">Y</td> <td style="vertical-align: top; padding: 5px;">Peptides, volume 5, no. 4, 1984, Ankho International Inc., (US), D.B. Richards et al.: "Effect of <math>\alpha</math>-MSH 11-13 (lysine-proline-valine) on fever in the rabbit", pages 815-817 see the whole document --</td> <td style="vertical-align: top; text-align: center; padding: 5px;">1-7</td> </tr> <tr> <td style="vertical-align: top; padding: 5px;">X,P</td> <td style="vertical-align: top; padding: 5px;">The Journal of Immunology, volume 137, no. 7, 1 October 1986, American Association of Immunologists, (US), J.G. Cannon et al.: "<math>\alpha</math> Melanocyte stimulating hormone inhibits immunostimulatory and inflammatory actions of interleukin 1", pages 2232-2236 see the whole document -----</td> <td style="vertical-align: top; text-align: center; padding: 5px;">1-7</td> </tr> </table>			Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>	X,Y	J. Rheumatol., volume 12, no. 5, October 1985, C.W. Denko et al.: "Effects of peptide hormones in urate crystal inflammation", pages 971-975 see the whole document --	1-7	Y	Peptides, volume 5, no. 4, 1984, Ankho International Inc., (US), D.B. Richards et al.: "Effect of $\alpha$ -MSH 11-13 (lysine-proline-valine) on fever in the rabbit", pages 815-817 see the whole document --	1-7	X,P	The Journal of Immunology, volume 137, no. 7, 1 October 1986, American Association of Immunologists, (US), J.G. Cannon et al.: " $\alpha$ Melanocyte stimulating hormone inhibits immunostimulatory and inflammatory actions of interleukin 1", pages 2232-2236 see the whole document -----	1-7
Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>												
X,Y	J. Rheumatol., volume 12, no. 5, October 1985, C.W. Denko et al.: "Effects of peptide hormones in urate crystal inflammation", pages 971-975 see the whole document --	1-7												
Y	Peptides, volume 5, no. 4, 1984, Ankho International Inc., (US), D.B. Richards et al.: "Effect of $\alpha$ -MSH 11-13 (lysine-proline-valine) on fever in the rabbit", pages 815-817 see the whole document --	1-7												
X,P	The Journal of Immunology, volume 137, no. 7, 1 October 1986, American Association of Immunologists, (US), J.G. Cannon et al.: " $\alpha$ Melanocyte stimulating hormone inhibits immunostimulatory and inflammatory actions of interleukin 1", pages 2232-2236 see the whole document -----	1-7												
<sup>10</sup> Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed		"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "d" document member of the same patent family												
<b>IV. CERTIFICATION</b> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; border-bottom: 1px solid black; padding: 5px;">           Date of the Actual Completion of the International Search            12th February 1988         </td> <td style="width: 50%; border-bottom: 1px solid black; padding: 5px;">           Date of Mailing of this International Search Report            15 APR 1988         </td> </tr> <tr> <td style="border-bottom: 1px solid black; padding: 5px;">           International Searching Authority            EUROPEAN PATENT OFFICE         </td> <td style="border-bottom: 1px solid black; padding: 5px;">           Signature of Authorized Officer             P.C.G. VAN DER PUTTEN         </td> </tr> </table>			Date of the Actual Completion of the International Search 12th February 1988	Date of Mailing of this International Search Report 15 APR 1988	International Searching Authority EUROPEAN PATENT OFFICE	Signature of Authorized Officer P.C.G. VAN DER PUTTEN								
Date of the Actual Completion of the International Search 12th February 1988	Date of Mailing of this International Search Report 15 APR 1988													
International Searching Authority EUROPEAN PATENT OFFICE	Signature of Authorized Officer P.C.G. VAN DER PUTTEN													